

# Lattice HQET Calculation of the Isgur-Wise Function

Joseph Christensen\*, Terrence Draper and Craig McNeile<sup>† a ‡</sup>

<sup>a</sup>Department of Physics and Astronomy, University of Kentucky, Lexington, KY 40506

We calculate the Isgur-Wise function on the lattice, simulating the light quark with the Wilson action and the heavy quark with a direct lattice implementation of the heavy-quark effective theory. Improved smearing functions produced by a variational technique, MOST, are used to reduce the statistical errors and to minimize excited-state contamination of the ground-state signal. Calculating the required matching factors, we obtain  $\xi'(1) = -0.64(13)$  for the slope of the Isgur-Wise function in continuum-HQET in the  $\overline{\text{MS}}$  scheme at a scale of 4.0 GeV.

## 1. The Tadpole-Improved Simulation

The Isgur-Wise function is the form factor of a heavy-light meson in which the heavy quark is taken to be much heavier than the energy scale,  $m_Q \gg \Lambda_{\text{QCD}}$ . This calculation adds perturbative corrections to the simulation results of Draper & McNeile [1]. The Isgur-Wise function is calculated using the action first suggested by Mandula & Ogilvie [2]:

$$iS = \sum_x \left\{ v_0 \left[ \psi^\dagger(x) \psi(x) - \psi^\dagger(x) \frac{U_t(x)}{u_0} \psi(x+\hat{t}) \right] + \sum_{j=1}^3 \frac{-iv_j}{2} \left[ \psi^\dagger(x) \frac{U_j(x)}{u_0} \psi(x+\hat{j}) - \psi^\dagger(x) \frac{U_j^\dagger(x-\hat{j})}{u_0} \psi(x-\hat{j}) \right] \right\}$$

This leads to the evolution equation:

$$G(x+\hat{t}) = \frac{U_t^\dagger(x)}{u_0} \left\{ G(x) - \sum_{j=1}^3 \frac{i\tilde{v}_j}{2} \left[ \frac{U_j(x)}{u_0} G(x+\hat{j}) - \frac{U_j^\dagger(x-\hat{j})}{u_0} G(x-\hat{j}) \right] \right\}$$

\*Presented by J. Christensen at Lattice '97, Edinburgh, Scotland.

<sup>†</sup>Currently at Department of Physics, University of Utah, Salt Lake City, UT 84112.

<sup>‡</sup>This work is supported in part by the U.S. Department of Energy under grant numbers DE-FG05-84ER40154 and DE-FC02-91ER75661, and by the University of Kentucky Center for Computational Sciences. The computations were carried out at NERSC.

where  $\tilde{v}_j = \frac{v_j}{v_0}$  and  $G(\vec{x}, t=0) = \frac{1}{v_0} f(\vec{x})$ . Better results can be obtained when a smeared source is used in place of a point source. The smearing function,  $f(\vec{x})$ , was calculated from a static simulation ( $v_j = 0$ ) *via* the smearing technique MOST (Maximal Operator Smearing Technique [3]).

## 2. The Isgur-Wise Function

The Isgur-Wise function was extracted from the lattice simulation as a ratio of three-point functions which was suggested by Mandula & Ogilvie [2].  $|\xi_{\text{unren}}^{\text{lat}}(v \cdot v')|^2$  is the large  $\Delta t$  limit of

$$\frac{4v_0 v'_0}{(v_0 + v'_0)^2} \frac{C_3^{vv'}(\Delta t) C_3^{v'v}(\Delta t)}{C_3^{vv}(\Delta t) C_3^{v'v'}(\Delta t)}$$

where  $\Delta t$  is the time separation between the current operator and each  $B$ -meson interpolating field.

Draper & McNeile have presented [1] the non-tadpole-improved unrenormalized slope of the lattice Isgur-Wise function to demonstrate the efficacy of the computational techniques.

## 3. Tadpole Improvement for HQET

Tadpole improvement grew from the observation that lattice links,  $U$ , have mean field value,  $u_0 \neq 1$ . Therefore, it is better to use an action written as a function of  $(U/u_0)$ . In the Wilson action, each link has a coefficient  $\kappa$ ;  $u_0$  can be paired with  $\kappa$  to easily tadpole improve *a posteriori* any previous non-tadpole-improved calculation.

In the HQET, there is no common coefficient (analogous to  $\kappa$ ) for both  $U_t$  and  $U_j$ . Correspondence between tadpole-improved and non-tadpole-improved HQET actions with  $v^2 = 1$  cannot be made *via* a simple rescaling of parameters, as is done for the Wilson action with  $\kappa$ .

Fortunately, the evolution equation can be written (as noticed by Mandula & Ogilvie [4]) such that the  $u_0$  is grouped with  $\tilde{v}_j$ . Thus, tadpole-improved Monte-Carlo data can be *constructed* from the non-tadpole-improved data by replacing  $v^{\text{nt}} \rightarrow v^{\text{tad}}$  and by including two overall multiplicative factors:

$$G^{\text{tad}}(t; \tilde{v}^{\text{tad}}, v_0^{\text{tad}}) = u_0^{-t} \frac{v_0^{\text{nt}}}{v_0^{\text{tad}}} G^{\text{nt}}(t; \tilde{v}^{\text{nt}}, v_0^{\text{nt}}) \quad (1)$$

In addition to the multiplicative factors  $u_0^{-t}$  and  $v_0^{\text{nt}}/v_0^{\text{tad}}$ , the tadpole-improvement of the simulation requires adjusting the velocity according to  $\tilde{v}^{\text{tad}} = u_0 \tilde{v}^{\text{nt}}$ , subject to  $(v^{\text{tad}})^2 = 1$  and  $(v^{\text{nt}})^2 = 1$ . Thus,

$$\begin{aligned} v_0^{\text{tad}} &= v_0^{\text{nt}} [1 + (1 - u_0^2)(v_j^{\text{nt}})^2]^{-1/2} \\ v_j^{\text{tad}} &= u_0 v_j^{\text{nt}} [1 + (1 - u_0^2)(v_j^{\text{nt}})^2]^{-1/2} \end{aligned}$$

#### 4. Tadpole-Improved Renormalization

By comparing the unrenormalized propagator

$$\left[ v_0^b \left( \frac{e^{ik_4}}{u_0} - 1 \right) + \frac{v_z^b}{u_0} \sin(k_z) + M_0^b - \Sigma(k, v) \right]^{-1}$$

to the renormalized propagator

$$iH(k, v) = Z_Q [v_0^r(ik_4) + v_z^r(k_z) + M^r]^{-1}$$

at  $O(k^2)$  and using  $(v^r)^2 = (v^b)^2 = 1$ , the perturbative renormalizations can be obtained. Aglietti [5] has done this for a different non-tadpole-improved action, for the special case  $\vec{v} = v_z \hat{z}$ .

With momentum shift,  $p \rightarrow p' = \langle p_4 + i \ln(u_0), \vec{p} \rangle$  [6], with  $\frac{1}{u_0} \exp(ik_4) = \exp(i(k_4 + i \ln(u_0)))$  and with  $X_\mu \equiv \frac{\partial}{\partial p_\mu} \Sigma(p)|_{p=0}$ , the tadpole-improved perturbative renormalizations are found to be

$$\begin{aligned} \delta M &= -\Sigma(0) - v_0 \ln(u_0) \\ \delta Z_Q &= Z_Q - 1 = -iv_0 X_4 - u_0 \sum_{j=1}^3 v_j X_j \end{aligned}$$

$$\delta \frac{v_i}{u_0} = -iv_0 \frac{v_i}{u_0} X_4 - (1 + v_i^2) X_i - v_i \sum_{j \neq i} v_j X_j$$

$$\delta v_0 = -i \sum_{j=1}^3 v_j^2 X_4 - u_0 v_0 \sum_{j=1}^3 v_j X_j$$

$u_0$  is the perturbative expansion and <sup>4</sup>

$$\begin{aligned} v_j^{r, \text{tad}} &= v_j^{b, \text{tad}} Z_{v_j}^{\text{tad}} \quad , \quad v_0^{r, \text{tad}} = v_0^{b, \text{tad}} Z_{v_0}^{\text{tad}} \\ Z_{v_j}^{\text{tad}} &\equiv \frac{1}{u_0} \left( 1 + \frac{\delta \frac{v_j}{u_0}}{\frac{v_j}{u_0}} \right) \quad , \quad Z_{v_0}^{\text{tad}} \equiv 1 + \frac{\delta v_0}{v_0} \end{aligned}$$

If one fits to  $\exp\{-t\}$  rather than  $\exp\{-(t+1)\}$ , the tadpole-improved wave-function renormalization is reduced to  $\delta Z'_Q = \delta Z_Q + (\delta M^{\text{tad}} + v_0 \ln(u_0))/v_0$ . Thus the  $\ln(u_0)$  term cancels explicitly and, as in the static case [6], tadpole-improvement has no effect on  $\delta Z'_Q$ , to order  $\alpha$ .

#### 5. Perturbative Renormalizations

We will present our computations of the renormalization factors elsewhere, but include this comment: Although the tadpole-improved functions include factors of  $u_0|_{\text{pt}}$  [7], these effects are higher order in  $\alpha$  and are dropped. Only the velocity renormalization is explicitly affected by the perturbative expansion of  $u_0$ :

$$Z_{v_j}^{\text{tad}} = \left( 1 + \frac{\delta \frac{v_j}{u_0}}{\frac{v_j}{u_0}} - \frac{g^2 C_F}{16\pi^2} (-\pi^2) \right)$$

The perturbative renormalizations favor a scale of  $q^* a = 1.9(1)$  for  $\alpha$ , which yields  $\alpha \approx 0.19(1)$ .

#### 6. Velocity Renormalization

Mandula & Ogilvie [4] consider the perturbative velocity renormalization expanded in orders of  $\tilde{v}$ . Our numbers for the velocity renormalization agree with theirs.

Another option is to consider, as did both Mandula & Ogilvie and Hashimoto & Matsufuru [8], the non-perturbative renormalization of the velocity. From Hashimoto's & Matsufuru's graph, we estimate their  $Z_{v, \text{np}}^{\text{tad}} \approx 1.05(5)$ . From Mandula & Ogilvie's result, we notice that  $Z_{\tilde{v}, \text{np}}^{\text{nt}} =$

<sup>4</sup>Note:  $(v^2 = 1) \Rightarrow \left( v_0 \delta v_0 = u_0^2 \sum_j \frac{v_j}{u_0} \frac{v_j}{\delta u_0} \right)$ .

$u_0 \times 1.01(1)$ . This is very close to the effect of tadpole-improving, and implies  $Z_v^{\text{tad}} = 1.01(1)$ .

We therefore use  $Z_v^{\text{tad}} \approx 1$  as the non-perturbative velocity renormalization in our calculation to renormalize the slope of the Isgur-Wise function.

## 7. Renormalization of $\xi'(v \cdot v')$

We claim that we can convert our Monte-Carlo data into tadpole-improved results and can calculate a renormalized tadpole-improved slope for the Isgur-Wise function.

We use the notation  $Z'_\xi = 1 + \delta Z'_\xi$  for the renormalization of the Isgur-Wise function, with  $\delta Z'_\xi$ :

$$\frac{g^2}{12\pi^2} [2(1 - v \cdot v' r(v \cdot v')) \ln(\mu a)^2 - f'(v, v')]$$

with  $r(v \cdot v')$  defined in [9] and primes on  $Z$  and  $f$  to indicate the “reduced value.”

For simplicity, we use the local current, which is not conserved on the lattice;  $Z'_\xi(1) \neq 1$ . However, the construction in §2 guarantees that the extracted renormalized Isgur-Wise function is properly normalized,  $\xi_{\text{ren}}(1) = 1$ .

## 8. Conclusions

After renormalization of our tadpole-improved results, we obtain  $\xi'_{\text{ren}}(1) = -0.64(13)$  for the slope of the Isgur-Wise function in continuum HQET in the  $\overline{\text{MS}}$  scheme at a scale of 4.0 GeV. Without renormalization, the slope is  $\xi'_{\text{unren}}(1) = -0.56(13)$ . Without tadpole-improvement, the slope is  $\xi'_{\text{unren}}^{\text{nt}}(1) = -0.43(10)$ .

We found that the tadpole-improved action (and therefore the tadpole-improved data) cannot be obtained from the non-tadpole-improved action (or data) by a simple rescaling of any parameter. However, the form of the evolution equation allows the *construction* of the tadpole-improved Monte-Carlo data from the non-tadpole-improved data as described in §3. After tadpole improvement, non-perturbative corrections to the velocity are negligible. Furthermore, tadpole improvement greatly reduces the perturbative corrections to the slope of the Isgur-Wise function.

### Unrenormalized Isgur-Wise Slope

Not Tadpole Improved				
$\Delta t$	0.154	0.155	0.156	$\kappa_c$
2	$0.38_{-1}^{+1}$	$0.38_{-1}^{+1}$	$0.38_{-1}^{+1}$	$0.39_{-1}^{+1}$
3	$0.42_{-2}^{+2}$	$0.41_{-2}^{+2}$	$0.41_{-2}^{+2}$	$0.41_{-2}^{+2}$
4	$0.50_{-9}^{+8}$	$0.48_{-9}^{+8}$	$0.45_{-10}^{+9}$	$0.43_{-10}^{+10}$

### Tadpole Improved

$\Delta t$	0.154	0.155	0.156	$\kappa_c$
2	$0.49_{-1}^{+1}$	$0.50_{-1}^{+1}$	$0.50_{-1}^{+1}$	$0.50_{-1}^{+1}$
3	$0.55_{-2}^{+3}$	$0.54_{-2}^{+3}$	$0.54_{-2}^{+3}$	$0.53_{-3}^{+3}$
4	$0.65_{-12}^{+10}$	$0.63_{-12}^{+11}$	$0.59_{-13}^{+12}$	$0.57_{-13}^{+13}$

### Renormalized Isgur-Wise Slope

Tadpole Improved				
$\Delta t$	0.154	0.155	0.156	$\kappa_c$
2	$0.57_{-1}^{+1}$	$0.57_{-1}^{+1}$	$0.58_{-1}^{+1}$	$0.58_{-1}^{+1}$
3	$0.62_{-2}^{+3}$	$0.62_{-2}^{+3}$	$0.61_{-2}^{+3}$	$0.61_{-3}^{+3}$
4	$0.72_{-11}^{+12}$	$0.71_{-11}^{+11}$	$0.67_{-11}^{+13}$	$0.64_{-13}^{+13}$

Table 1

The negative of the slope at the normalization point,  $\xi'(1)$ , from both the unrenormalized and the renormalized ratio of three-point functions. This ratio gives the (un)-renormalized Isgur-Wise function  $\xi(v \cdot v')$  at asymptotically-large times  $\Delta t$ .

## REFERENCES

1. T. Draper and C. McNeile, Nucl. Phys. B (Proc. Suppl.) **47**, 429 (1996).
2. J.E. Mandula and M.C. Ogilvie, Phys. Rev. D **45**, R2183 (1992).
3. T. Draper and C. McNeile, Nucl. Phys. B (Proc. Suppl.) **34**, 453 (1994).
4. J.E. Mandula and M.C. Ogilvie, [hep-lat/9602004](#), [hep-lat/9703020](#).
5. U. Aglietti, Nucl. Phys. **B421**, 191 (1994).
6. C. Bernard, Nucl. Phys. B (Proc. Suppl.) **34**, 47 (1994).
7. G. P. Lepage and P. B. Mackenzie, Phys. Rev. D **48**, 2250 (1993).
8. S. Hashimoto and H. Matsufuru, Phys. Rev. D **54**, 4578 (1996).
9. A. F. Falk, H. Georgi, B. Grinstein, and M. B. Wise, Nucl. Phys. **B343**, 1 (1990).